


1. Consider the following reaction in a closed reaction flask:  $2A(g) + 3B(g) \rightarrow A_2B_3(g)$  If 1.20 atm of gas A is allowed to react with 1.20 atm of gas B, and the reaction goes to completion at constant temperature and volume, what is the total pressure (in atm) in the reaction flask at the end of the reaction?

a 0.4 atm

b 0.8 atm

c 1.2 atm

d 2.4 atm



$$1.2 \text{ atm A} \cdot \frac{1 \text{ mol } A_2B_3}{2 \text{ mol A}} = 0.6 \text{ atm } A_2B_3$$


$$1.2 \text{ atm B} \cdot \frac{1 \text{ mol } A_2B_3}{3 \text{ mol B}} = 0.4 \text{ atm } A_2B_3^*$$

$$0.6 \text{ atm } A_2B_3 - 0.4 \text{ atm } A_2B_3 = 0.2 \text{ atm } A_2B_3$$

$$0.2 \text{ atm } A_2B_3 \cdot \frac{2 \text{ mol A}}{1 \text{ mol } A_2B_3} = 0.4 \text{ atm A in excess}$$

$$P_f = \sum P_{A_2B_3} + \sum P_{A(\text{ex})} = 0.4 \text{ atm } A_2B_3 + 0.4 \text{ atm A} = 0.8 \text{ atm gas}$$

2. A mixture of  $Xe(g)$  and  $O_2(g)$ , formed by the complete decomposition of  $XeO_4(g)$ , is collected over water at  $34^\circ C$  at a total pressure of 760 mmHg. If the vapor pressure of water is 40 mmHg at  $34^\circ C$ , what is the partial pressure of  $O_2(g)$ ? If  $O_2(g)$  is isolated in a 250 mL container at the same temperature, how many grams of  $O_2(g)$  is produced?



$$\delta P_{H_2O} = 40 \text{ mmHg}$$

$$P_f = 760 \text{ mmHg}$$

$$P_f = \delta P_{O_2} + \delta P_{Xe} + \delta P_{H_2O}$$

$$\delta P_{O_2} = 2x$$

$$= 2 \cdot (240 \text{ mmHg})$$

$$\delta P_{O_2} = 480 \text{ mmHg}$$

$$760 \text{ mmHg} = \delta P_{O_2} + \delta P_{Xe} + 40 \text{ mmHg}$$

$$x \text{ atm Xe} \cdot \frac{2 \text{ mol } O_2}{1 \text{ mol Xe}} = 2x \text{ atm } O_2$$

$$760 \text{ mmHg} = \frac{2x + x}{3} + 40 \text{ mmHg}$$

$$\frac{720 \text{ mmHg}}{3} = \frac{3x}{3}$$

$$x = 240 \text{ mmHg} = \delta P_{Xe}$$

$$P_{O_2} = 480 \text{ mmHg}$$

$$\frac{480 \text{ mmHg}}{760 \text{ mmHg}} \cdot 1 \text{ atm} = .632 \text{ atm}$$

$$V_{O_2} = 250 \text{ mL}$$

$$T_{O_2} = 34^\circ C$$



$$m = ?$$

$$PV = nRT$$

$$d = \frac{MP}{RT}$$

$$M = \frac{mRT}{PV}$$

molar mass  
gram



$$\frac{(.632 \text{ atm})(.25 \text{ L})}{.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} \cdot 307.15 \text{ K}} = \frac{m \cdot (10.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}})(307.15 \text{ K})}{(.632 \text{ atm})(.25 \text{ L})}$$

$$m = 0.20 \text{ g O}_2$$

3. In an experiment, 25.0 ml of a gas with a pressure of 1.00 atm is contained in a balloon at 25.00°C. The balloon's temperature is adjusted until the pressure is 0.75 atm at a volume of 31.1 ml. What is the final temperature of the gas under the new conditions?

$$\frac{P_1 V_1}{n_1 T_1} = \frac{P_2 V_2}{n_2 T_2} \quad T_2 \cdot \frac{(1 \text{ atm})(25 \text{ mL})}{298.15 \text{ K}} = \frac{.75 \text{ atm} \cdot 31.1 \text{ mL}}{T_2} \cdot \frac{298.15 \text{ K}}{1 \text{ atm} \cdot 25 \text{ mL}} \cdot T_2$$

-cancel out anything that stays the same

$$T_2 = \frac{.75 \text{ atm} \cdot 31.1 \text{ mL} \cdot 298.15 \text{ K}}{1 \text{ atm} \cdot 25 \text{ mL}}$$

$$T_2 = 278.17 \text{ K} - 273.15 = 5.02^\circ\text{C}$$

4. If 1000. g of boiling water (at 100°C) was placed in an 1800. g cast iron skillet initially at 25°C, and the final equilibrium temperature of the water and the skillet was 88°C, estimate the specific heat capacity of the skillet. Assume this is a closed system and that the specific heat capacity of water is 4.184 J/°C·g.

$$m_w = 1000 \text{ g}$$

$$T_{i,w} = 100^\circ\text{C}$$

$$m_p = 1800 \text{ g}$$

$$T_{i,p} = 25^\circ\text{C}$$

$$c_w = 4.184 \text{ J/g}\cdot^\circ\text{C}$$

$$T_f = 88^\circ\text{C}$$

$$-q_{\text{water}} = q_{\text{pan}} \quad q = mc\Delta T \quad c_p$$

$$-(1000 \text{ g} \cdot 4.184 \text{ J/g}\cdot^\circ\text{C} \cdot (88^\circ\text{C} - 100^\circ\text{C})) = (1800 \text{ g} \cdot x \cdot (88^\circ\text{C} - 25^\circ\text{C}))$$

$$\frac{50208 \text{ J}}{113400 \text{ g}\cdot^\circ\text{C}} = \frac{113400 \text{ g}\cdot^\circ\text{C} \cdot x}{113400 \text{ g}\cdot^\circ\text{C}}$$

$$0.443 \text{ J/g}\cdot^\circ\text{C} = c_p$$

5. Which statement is incorrect regarding internal energy (U, E) and the first law of thermodynamics?

A) The first law of thermodynamics states that energy must be conserved. ✓  $\Delta E = q + w$


B) When the system gains heat and performs work, then  $\Delta\{U, E\}$  for the system must be positive.  
 $+q \quad -w \quad U = q + w \rightarrow +q < -w$

C) The first law of thermodynamics does not imply that heat can't be converted to work. ✓  $\Delta E = q + w$

D) When the system loses heat and performs work, then  $\Delta\{U, E\}$  for the system must be negative. ✓  
 $-q \quad -w \quad U = -q + -w = -U$

E) When its  $\Delta\{U, E\}$  increases, then the system must gain heat or have work performed on it, or both. ✓  
 $+q \quad +w$

6. Deterioration of buildings, bridges, and other structures through the rusting of iron costs millions of dollars a day. The enthalpy of formation of rust,  $\text{Fe}_2\text{O}_3(\text{s})$ , is  $-826.0 \text{ kJ/mol}$ . How much heat is released (in kJ) when  $0.500 \text{ kg}$  of Fe reacts with  $200. \text{ g}$  of  $\text{O}_2$ , forming  $\text{Fe}_2\text{O}_3(\text{s})$ ?



$$0.5 \text{ kg Fe} \cdot \frac{1000 \text{ g}}{1 \text{ kg}} \cdot \frac{1 \text{ mol Fe}}{55.85 \text{ g Fe}} \cdot \frac{1 \text{ mol Fe}_2\text{O}_3}{2 \text{ mol Fe}} = 4.476 \text{ mol Fe}_2\text{O}_3$$

$$200 \text{ g O}_2 \cdot \frac{1 \text{ mol O}_2}{32 \text{ g O}_2} \cdot \frac{1 \text{ mol Fe}_2\text{O}_3}{\frac{3}{2} \text{ mol O}_2} = 4.167 \text{ mol Fe}_2\text{O}_3$$

amount heat

$$\Delta H_f = \frac{q}{n}$$

$$-826.0 \text{ kJ/mol} = \frac{q}{4.167 \text{ mol Fe}_2\text{O}_3}$$


$$= -3441.67 \text{ kJ}$$

heat is release

**3440 kJ heat released**

**exothermic** When  $50.0 \text{ ml}$  of  $0.200 \text{ M AgNO}_3$  and  $50.0 \text{ ml}$  of  $0.100 \text{ M CaCl}_2$ , both at  $25.0^\circ\text{C}$ , are reacted in a coffee-cup calorimeter, the temperature of the reacting mixture increases to  $26.0^\circ\text{C}$ . Calculate  $\Delta H$  in kJ per mole of  $\text{AgCl}$  produced. Assume the density of the solution is  $1.05 \text{ g/ml}$  and the specific heat capacity of the solution  $4.20 \text{ J/g}^\circ\text{C}$ .

$$\frac{q}{n} = -\Delta H$$



$$\frac{.2 \text{ mol AgNO}_3 \cdot .050 \cancel{\text{L}} \cdot 2 \text{ mol AgCl}}{2 \text{ mol AgNO}_3} = 0.01 \text{ mol AgCl}$$

$\Delta H = \frac{q}{n}$   
amount of heat / amount AgCl produced

$$\frac{.100 \text{ mol CaCl}_2 \cdot .05 \cancel{\text{L}} \cdot 2 \text{ mol AgCl}}{1 \text{ mol CaCl}_2} = .01 \text{ mol AgCl} = n$$

$$\Delta H = \frac{441 \text{ J}}{.01 \text{ mol}}$$

$$= 44,100 \text{ J/mol}$$

**-44.1 kJ/mol**

$$q = mc\Delta T$$

(100 mL · 1.05 g/mL) · 4.20 J/g°C · 26°C - 25°C

$$q = 105 \text{ g} \cdot 4.2 \frac{\text{J}}{\text{g}^\circ\text{C}} \cdot 1^\circ\text{C} = 441 \text{ J} = q$$

↑ +q = gaining heat

8. A pure gold ring ( $C = 0.128 \text{ J/g}^\circ\text{C}$ ) and pure silver ring ( $C = 0.235 \text{ J/g}^\circ\text{C}$ ) have a total mass of 15.3g. The two rings are heated to 62.1°C and dropped into a 13.1mL of water ( $\rho = 1.00 \text{ g/mL}$  and  $C = 4.184 \text{ J/g}^\circ\text{C}$ ) at 20.9°C. When equilibrium is reached, the temperature of the water is 22.9°C. What was the mass of the gold ring?

$$\rightarrow m_g + m_s = 15.3 \text{ g}$$

$$q_{\text{system}} = -q_{\text{surroundings}} \quad m_s = 15.3 \text{ g} - m_g$$

$$-(q_s + q_g) = q_w$$

$$T_{i,gs} = 62.1^\circ\text{C}$$

$$T_{i,w} = 20.9^\circ\text{C}$$

$$T_f = 22.9^\circ\text{C}$$

$$C_g = .128 \text{ J/g}^\circ\text{C}$$

$$C_s = .235 \text{ J/g}^\circ\text{C}$$

$$+(m_g \cdot .128 \frac{\text{J}}{\text{g}^\circ\text{C}} (+39.2^\circ\text{C})) + (m_s \cdot .235 \frac{\text{J}}{\text{g}^\circ\text{C}} (+39.2^\circ\text{C})) = (13.1 \text{ mL} \cdot 1.0 \frac{\text{g}}{\text{mL}}) (4.184 \frac{\text{J}}{\text{g}^\circ\text{C}}) (2.0^\circ\text{C})$$

$$\underline{5.0176 m_g} + \underline{140.9436 m_s} - \underline{9.212 m} = 109.62 \text{ J} - 140$$

$$\frac{-4.1944 m_g}{-4.1944} = \frac{-31.3236 \text{ J}}{-4.1944}$$


$$m_g = 7.5 \text{ g}$$

9. Find the heat of formation of gaseous HCl



$$\Delta H = (-91.8 \text{ kJ}) - \frac{1}{2}$$

$$-\frac{1}{2} \cdot -91.8$$



$$\Delta H = -628.8 \text{ kJ} \cdot \frac{1}{2}$$





$$+ \frac{1}{2} \cdot -628.8$$



$$\Delta H = (-176.2 \text{ kJ}) -$$


$$+ (-176.2) - 1$$

$$2 - \frac{3}{2} = \frac{1}{2}$$



Flipped  $-92.3 \text{ kJ}$

10. Consider the reaction



in which 10.0 g of sucrose,  $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ , was burned in a bomb calorimeter with a heat capacity of  $7.50 \text{ kJ}/^\circ\text{C}$ . The temperature increase inside the calorimeter was found to be  $22.0^\circ\text{C}$ . What is the heat of this reaction per mole of sucrose?

$q \rightarrow \text{J}$   
 $\Delta H \rightarrow \text{kJ/mol}$



$$-q_{\text{sample}} = q_{\text{water}} + q_{\text{cal}}$$

$C_{\text{sucrose}} = 7.50 \text{ kJ}/^\circ\text{C} \cdot 22^\circ\text{C}$

$$10 \text{ g } \text{C}_{12}\text{H}_{22}\text{O}_{11} \cdot \frac{1 \text{ mol}}{342.3 \text{ g}} = 0.0292 \text{ mol}$$


$$\Delta H = \frac{165000 \text{ J}}{0.0292 \text{ mol}} = 5650 \text{ kJ/mol}$$

**- 5650 kJ/mol**

$$q = 7500 \text{ J}/^\circ\text{C} \cdot 22^\circ\text{C} = 165000 \text{ J}$$

$$\Delta H = q/n$$

11. For each of the following orbital shapes below, give the maximum number of electrons that can be accommodated in the orbitals that share the same principal quantum number, n, and angular quantum number, l.



A) 4, 2 B) 6, 2 C) 5, 3 D) 10, 6 E) 14, 10

$$m_l = \begin{matrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{matrix} \left\{ \begin{matrix} 5 \times 2e^- \\ = 10e^- \end{matrix} \right.$$

$d \rightarrow l=2$


$p \rightarrow l=1$

$m_l = -1, 0, 1 \rightarrow 3 \text{ orbitals} \times 2e^- = 6e^-$







12. Which of the following full sets of quantum numbers is incorrect?

- a) The  $e^-$  gained from  $\text{Br} \rightarrow (\text{Br})^-$ ;  $n=4, l=1, m_l=+1, m_s=-1/2$  ✓
- b) The outermost  $e^-$  in  $\text{Rb}$ ;  $n=5, l=0, m_l=0, m_s=+1/2$  ✓
- c) The 6<sup>th</sup>  $e^-$  in  $\text{O}$ ;  $n=2, l=1, m_l=0, m_s=+1/2$  ✗
- d) The 3<sup>rd</sup>  $e^-$  in  $\text{F}$ ;  $n=2, l=0, m_l=0, m_s=+1/2$  ✓
- e) The 8<sup>th</sup>  $e^-$  in  $\text{O}$ ;  $n=2, l=1, m_l=-1, m_s=-1/2$  ✓

n	l	$m_l$	$m_s$
energy level	shape	orientation	spin electron
low #	$s=0$	$\pm l$	$\pm 1/2$
	$p=1$	$p=-1, 0, 1$	
$d: n=r-1$	$d=2$	$d=-2 \rightarrow +2$	
$f: n=r-2$	$f=3$		



13. Which of the following electron configurations are correct?




a) I, III, V

b) II, IV


c) I, II, V, VI

d) II, III, IV, V




e) None



14. What is one correct set of quantum numbers for the third electron removed to form a cation of nickel? If there are multiple, give a range for each quantum number.



15. Which of these are in the correct increasing atomic size order?




16. Which of these are in the correct order for increasing  $IE_1$ .

- a) Cs < Xe < I ✗
- b) Kr < Ar < He ✓
- c) Rb < Ca < K ✗
- d) Sn < Sb < I ✗
- e) A and C
- f) B and D

inc. →  
↑ inc.  
Cations

IE = energy to remove  $e^-$   
electron affinity = energy to add one  $e^-$



17. If a light bulb consumes 218 J per second, and all of its energy is converted to 560 nm light, how many photons are produced per second?


218 J/s → ? = photons/sec       $\lambda = 560 \text{ nm}$

$$E = h\nu \quad \left. \begin{array}{l} \uparrow \\ \text{frequency} \end{array} \right\} \nu = \frac{c}{\lambda}$$

$$E = \frac{h \cdot c}{\lambda} \rightarrow \frac{6.626 \times 10^{-34} \text{ J}\cdot\text{s} \cdot 3 \cdot 10^8 \text{ m/s}}{560 \times 10^{-9} \text{ m}} = 3.5496 \times 10^{-19} \text{ J/photon}$$

$$\frac{1 \text{ photon}}{3.5496 \times 10^{-19} \text{ J}} \cdot \frac{218 \text{ J}}{\text{s}} = 6.14 \times 10^{20} \text{ photons/s}$$

18. Label the following ions paramagnetic or diamagnetic. → all  $e^-$  have pairs



19. Which ions are ranked correctly by decreasing size?

- ✓ I.  $\text{Si}^{2+} > \text{Ca}^{2+} > \text{Mg}^{2+}$    
 ✓ II.  $\text{S}^{2-} > \text{Cl}^- > \text{K}^+$    
 ✗ III.  $\text{Mg}^{2+} > \text{Na}^+ > \text{F}^-$    
 ✗ IV.  $\text{Ba}^{2+} > \text{Cs}^+ > \text{I}^-$

← inc.  
↓ inc.

✓ V.  $\text{P}^{3-} > \text{S}^{2-} > \text{Cl}^-$

isoelectronic

biggest ↓

$\text{Mg}^{2+} < \text{Na}^+ < \text{F}^-$

isoelectronic \*  
cations < anions

↑ anion >>  
 $\text{S}^{2-} > \text{Cl}^- > \text{K}^+$

- a) I, III, V
- b) II, IV
- c) I, II, V
- d) I, IV, V
- e) II, III, IV, V

