

- A. Sign your bubble sheet on the back at the bottom in ink.
- B. In pencil, write and encode in the spaces indicated:
- 1) Name (last name, first initial, middle initial)
 - 2) UF ID number
 - 3) **SKIP** Section number
- C. Under "special codes" code in the test ID numbers 4, 2.
- | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| 1 | 2 | 3 | ● | 5 | 6 | 7 | 8 | 9 | 0 |
| 1 | ● | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 0 |
- D. At the top right of your answer sheet, for "Test Form Code", encode B.
- A ● C D E
- E. 1) This test consists of 22 multiple choice questions. The test is counted out of 100 points, and there are 10 bonus points available.
- 2) The time allowed is 120 minutes.
 - 3) Raise your hand if you need more scratch paper or if you have a problem with your test. **DO NOT LEAVE YOUR SEAT UNLESS YOU ARE FINISHED WITH THE TEST.**
- F. KEEP YOUR BUBBLE SHEET COVERED AT ALL TIMES.**
- G. When you are finished:
- 1) Before turning in your test **check carefully for transcribing errors**. Any mistakes you leave in are there to stay.
 - 2) You must turn in your scantron to your discussion leader or exam proctor. Be prepared to show your picture I.D. with a legible signature.
 - 3) The answers will be posted in Canvas within one day after the exam.

University of Florida Honor Pledge:

On my honor, I have neither given nor received unauthorized aid doing this exam.

Signature: _____

Summary of Integration Formulas

- Fundamental Theorem of Calculus

$$\int_a^b F'(x) dx = F(b) - F(a)$$

- Fundamental Theorem of Line Integrals

$$\int_C \nabla f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$$

- Green's Theorem (circulation form)

$$\iint_D \text{curl } \vec{F} \cdot \hat{k} dA = \oint_C \vec{F} \cdot d\vec{r}$$

- Stokes' Theorem

$$\iint_S \text{curl } \vec{F} \cdot \hat{n} dS = \oint_C \vec{F} \cdot d\vec{r}$$

- Green's Theorem (flux form)

$$\iint_D \text{div } \vec{F} dA = \oint_C \vec{F} \cdot \hat{n} ds$$

- Divergence Theorem

$$\iiint_E \text{div } \vec{F} dV = \oiint_S \vec{F} \cdot \hat{n} dS$$

NOTE: Be sure to bubble the answers to questions 1–22 on your scantron.

Questions 1 – 22 are worth 5 points each.

1. Let $\vec{F}(x, y, z) = \langle x, -2yz, 3xz^2 \rangle$. Which of the following vectors is orthogonal to $\text{curl } \vec{F}$ at the point $(1, 1, 1)$?

- a. $\langle 2, -3, 0 \rangle$
- b. $\langle 2, 3, 0 \rangle$
- c. $\langle 3, -2, 5 \rangle$
- d. $\langle -3, -2, 1 \rangle$
- e. $\langle -3, 2, -1 \rangle$

2. Let $\vec{F} = \langle x^2 - y^2, -2xy + y \rangle$. Which of the following statements must be correct?

P. $\nabla \cdot \vec{F} = 0$.

Q. \vec{F} is conservative.

R. $\int_C \vec{F} \cdot d\vec{r} = 0$ for any smooth curve C .

- a. P and Q only
- b. Q only
- c. P and R only
- d. Q and R only
- e. P, Q, and R

3. Evaluate the line integral $\int_C 2xe^y ds$, where C is the line segment from $(0, 0)$ to $(3, 1)$.

- a. 6
- b. $6e$
- c. $6(e - 1)$
- d. $6\sqrt{10}(e - 1)$
- e. $6\sqrt{10}$

4. Calculate $\oint_C \frac{y}{2} dx$, where C is the counterclockwise oriented curve bounding the triangle with vertices $(0, 0)$, $(4, 0)$, and $(1, 3)$.

- a. -3
- b. -6
- c. 0
- d. 3
- e. 6

7. If f is a potential function of $\vec{F}(x, y) = \langle -y \sin(xy), -x \sin(xy) - 2y \rangle$ and $f(0, 0) = 3$, find $f(0, 2)$.

- a. 1
- b. 0
- c. -1
- d. -2
- e. -3

8. If $\vec{F} = \langle -x, 0, z \rangle$, which of the following must be correct?

- P. The flux of \vec{F} across the plane $z = 1$ is 0.
 - Q. The flux of \vec{F} across the plane $x = 1$ is 0.
 - R. The flux of \vec{F} across a unit sphere is 0.
-
- a. P only
 - b. Q only
 - c. R only
 - d. P and Q only
 - e. P, Q, and R

5. The surface S is parameterized by $\vec{r}(u, v) = \langle 2 \sin(v) \cos(u), 2 \sin(v) \sin(u), 2 \cos(v) \rangle$, $0 \leq u \leq 2\pi$ and $0 \leq v \leq \pi$. Which of the following statements is/are correct?

P. The surface S is the sphere centered at $(0, 0, 0)$ with radius 4.

Q. The vector $\vec{r}_u(P)$ is parallel to the tangent plane to S at the point P .

R. The area of the surface $S = \iint_D dA$, where $D = \{(u, v) \mid 0 \leq u \leq 2\pi, 0 \leq v \leq \pi\}$.

- a. P only
- b. Q only
- c. R only
- d. P and Q
- e. Q and R

6. Find the area of the surface S , where S is the part of the plane $2x + y + 2z = 10$ that lies inside the cylinder $x^2 + y^2 = 16$.

- a. 48π
- b. 18π
- c. 16π
- d. 12π
- e. 24π


9. Let $\vec{F}(x, y, z) = \langle x, y, -2xy \rangle$. Evaluate the line integral $\int_C \vec{F} \cdot d\vec{r}$, where C is the curve parameterized by $\vec{r}(t) = \langle \cos t, \sin t, 2t \rangle$, $0 \leq t \leq \frac{\pi}{2}$.

- a. $-\pi$
- b. π
- c. 2
- d. -2
- e. 0

10. If the surface S is parameterized by $\vec{r}(u, v) = \langle u, v \cos(2u), v \sin(2u) \rangle$, find an equation of the tangent plane to S at the point $(\pi, 1, 0)$.

- a. $-2x + z + \pi = 0$
- b. $-2x + z + 2\pi = 0$
- c. $2x + z - 2\pi = 0$
- d. $2y + z - \pi = 0$
- e. $-2y + z + 2\pi = 0$

11. Which of the following vector fields has the graph below?



- a. $\vec{F}_1(x, y) = -y\hat{i} + x\hat{j}$
 b. $\vec{F}_2(x, y) = \hat{i} + \hat{j}$
 c. $\vec{F}_3(x, y) = \hat{i} + x\hat{j}$
 d. $\vec{F}_4(x, y) = x\hat{i} + y\hat{j}$
 e. $\vec{F}_5(x, y) = y\hat{i} + \hat{j}$

12. Let $\vec{F}(x, y, z) = \langle 2y^3, 1, e^z \rangle$. Find the circulation of \vec{F} along C , where C is the curve of intersection of the plane $y + 2z = 3$ and the cylinder $x^2 + y^2 = 4$. (Orient C to be counterclockwise when viewed from above.) By Stoke's Theorem,

$$\int_C \vec{F} \cdot d\vec{r} =$$

- a. $\int_0^{2\pi} \int_0^2 -6r^3 \sin^2 \theta \, dr \, d\theta$
 b. $\int_0^{2\pi} \int_0^2 -12r^3 \sin^2 \theta \, dr \, d\theta$
 c. $\int_0^{2\pi} \int_0^2 12r^2 \cos^2 \theta \, dr \, d\theta$
 d. $\int_0^{2\pi} \int_0^2 12r^3 \cos^2 \theta \, dr \, d\theta$
 e. $\int_0^{2\pi} \int_0^2 -6r^2 \sin^2 \theta \, dr \, d\theta$

15. Find the work done by the force $\vec{F} = \langle 3x^2, 3y^2 \rangle$ in moving a particle along the parametric curve $\vec{r}(t) = \langle t \cos t, t \sin t \rangle$, $0 \leq t \leq 2\pi$.

Hint: Is \vec{F} conservative?

- a. $8\pi^3$
- b. 8π
- c. 2π
- d. $2\pi^3$
- e. 0

16. Calculate $\iint_S (\nabla \times \vec{F}) \cdot \hat{n} \, dS$, where $\vec{F} = \langle y, -x, e^{xyz} \rangle$ and S is the part of paraboloid $z = 3x^2 + 3y^2$, $0 \leq z \leq 6$, oriented downward.

- a. -2π
- b. -4π
- c. 0
- d. 2π
- e. 4π

13. Which of the following is correct?

- a. If \vec{F} is conservative, then $\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r}$ for any two smooth curves C_1 and C_2 .
- b. If \vec{F} is conservative, then $\iint_S \vec{F} \cdot d\vec{S} = 0$.
- c. If \vec{F} is conservative, then $\operatorname{div} \vec{F} = 0$.
- d. If D is a simply connected planar region, then the area of D is $\oint_{\partial D} \frac{y}{2} dx - \frac{x}{2} dy$, where ∂D is oriented counterclockwise.
- e. If $\vec{F} = \langle x, -2y, z \rangle$, then $\oiint_S \vec{F} \cdot d\vec{S} = 0$, where S is a unit sphere.
-

14. Set up a double integral for the surface integral $\iint_S (z + 1) dS$, where S is the part of the paraboloid $z = x^2 + y^2 - 1$, $-1 \leq z \leq 5$.

- a. $\int_0^{2\pi} \int_0^1 r^3 \sqrt{4r^2 + 1} dr d\theta$
- b. $\int_0^{2\pi} \int_0^1 r^2 dr d\theta$
- c. $\int_0^{2\pi} \int_0^{\sqrt{6}} r^2 dr d\theta$
- d. $\int_0^{2\pi} \int_0^{\sqrt{6}} r^3 \sqrt{4r^2 + 1} dr d\theta$
- e. $\int_0^{2\pi} \int_0^{\sqrt{6}} r^2 \sqrt{4r^2 + 1} dr d\theta$

17. Let $\vec{F} = \langle x^3 + y^3, y^3 + z^3, z^3 + x^3 \rangle$ and let S be the sphere centered at $(0, 0, 1)$ with radius 1. Then the flux of \vec{F} across the surface S is

a. $\int_0^{2\pi} \int_0^{\pi} \int_0^1 3\rho^4 \sin \phi \, d\rho \, d\phi \, d\theta$

b. $\int_0^{2\pi} \int_0^{\pi/2} \int_0^{\cos \phi} 3\rho^4 \sin \phi \, d\rho \, d\phi \, d\theta$

c. $\int_0^{2\pi} \int_0^{\pi/2} \int_0^{\cos \phi} 3\rho^2 \, d\rho \, d\phi \, d\theta$

d. $\int_0^{2\pi} \int_0^{\pi/2} \int_0^{2 \cos \phi} 3\rho^4 \sin \phi \, d\rho \, d\phi \, d\theta$

e. $\int_0^{2\pi} \int_0^{\pi} \int_0^{2 \cos \phi} 3\rho^2 \, d\rho \, d\phi \, d\theta$

18. Which of the following regions is/are simply connected?

$$D_1 = \{ (x, y) \mid 1 < x^2 + y^2 < 9 \text{ and } x > 0 \}$$

$$D_2 = \{ (x, y) \mid x > 0 \text{ and } y > 0 \}$$

$$D_3 = \{ (x, y) \mid y \neq 1 \}$$

- a. D_1 only
b. D_2 only
c. D_3 only
d. D_1 and D_2
e. D_2 and D_3

19. Let D be the region bounded by $y = \sqrt{2x - x^2}$ and the x -axis and let ∂D be its boundary curve oriented positively. Then

$$\int_{\partial D} -x^2 y \, dx + xy^2 \, dy =$$

a. $\int_0^\pi \int_0^{2 \cos \theta} -r^2 \, dr \, d\theta$

b. $\int_0^{\pi/2} \int_0^{2 \cos \theta} r^3 \, dr \, d\theta$

c. $\int_0^{\pi/2} \int_0^{2 \cos \theta} -r^3 \, dr \, d\theta$

d. $\int_0^\pi \int_0^{2 \cos \theta} r^3 \, dr \, d\theta$

e. $\int_0^{\pi/2} \int_0^{2 \cos \theta} r^2 \, dr \, d\theta$

20. Let $\vec{F}(x, y, z) = (3x - y)\hat{i} + (3y - z)\hat{j} + (3z - x)\hat{k}$ and let S be the surface of the solid bounded by $x = 0$, $x = 2$, $y = 0$, $y = 2$, $z = 0$, and $z = 2$. Find the flux of \vec{F} across S .

a. 54

b. 108

c. 36

d. 72

e. 18

21. Let $\vec{F}(x, y, z) = \langle \cos(\sqrt{x^2 + y^2 + z^2}), e^{\sqrt{xyz}}, \sin^3(x) \rangle$. Find $\text{div}(\text{curl } \vec{F})$ at the point $(1, 1, 1)$.

- a. 0
- b. e
- c. $\frac{e}{2}$
- d. \sqrt{e}
- e. 1

22. Let $\vec{F}(x, y) = \langle x^3 - y^2, x^2 - y^3 \rangle$. Let D be the region bounded by $y = x^2$, $y = 0$ and $x = 1$, and ∂D be its boundary curve oriented positively. Then the flux of \vec{F} across the boundary curve $\partial D = \oint_{\partial D} \vec{F} \cdot \hat{n} \, ds =$

- a. $\int_0^1 \int_0^{x^2} (2x + 2y) \, dy \, dx$
- b. $\int_0^1 \int_{x^2}^1 (2x + 2y) \, dy \, dx$
- c. $\int_0^1 \int_0^{x^2} (3x^2 + 3y^2) \, dy \, dx$
- d. $\int_0^1 \int_{x^2}^1 (3x^2 - 3y^2) \, dy \, dx$
- e. $\int_0^1 \int_0^{x^2} (3x^2 - 3y^2) \, dy \, dx$